

Week 05 Quiz: Sampling Distribution

Problem 1

The daily revenue at a university snack bar has been recorded for the past five years. Records indicate that the mean daily revenue is \$1500, and the standard deviation is \$500. The distribution is skewed to the right due to several high-volume days (football game days). Suppose that 100 days are randomly selected and the average daily revenue computed. Which of the following describes the sampling distribution of the sample mean?

- A). normally distributed with a mean of \$1500 and a standard deviation of \$500
- B). normally distributed with a mean of \$1500 and a standard deviation of \$50
- C). normally distributed with a mean of \$150 and a standard deviation of \$50
- D). skewed to the right with a mean of \$1500 and a standard deviation of \$500

Answer: B) standard error $s_{\bar{x}} = 500/\sqrt{100}=50$, $\bar{X} \sim N(1500, 50)$

Problem 2.

Suppose students' ages follow a skewed right distribution with a mean of 23 years old and a standard deviation of 4 years. If we randomly sample 200 students, which of the following statements about the sampling distribution of the sample mean age is incorrect?

- A) The mean of sampling distribution is approximately 23 years old.
- B) The standard deviation of the sampling distribution is equal to 4 years.
- C) The shape of the sampling distribution is approximately normal

Answer: C). This is based on the central limit theorem. $s_{\bar{x}} = 4/\sqrt{200}=0.283$, $\bar{X} \sim N(23, 0.283)$

Problem 3

A random sample of $n = 600$ measurements is drawn from a binomial population with probability of success .08. What is the sampling distribution of the sample proportion, \hat{p} .

- A) $N(.08; .011)$
- B) $N(.92; .003)$
- C) $N(.08; .003)$
- D) $N(.92; .011)$

Answer: A) $s_{\hat{p}} = \sqrt{0.08(1 - 0.08)/600} = 0.0111$

Problem 4

You form a distribution of the means of all samples of size 36 drawn from an infinite population that is skewed to the left. The population from which the samples are drawn has a mean of 50 and a standard deviation of 12 . Which one of the following statements is true of this distribution?

- (a) $\mu_{\bar{x}} = 50, \sigma_{\bar{x}} = 12$, the sampling distribution is skewed somewhat to the left.
- (b) $\mu_{\bar{x}} = 50, \sigma_{\bar{x}} = 2$, the sampling distribution is skewed somewhat to the left.
- (c) $\mu_{\bar{x}} = 50, \sigma_{\bar{x}} = 12$, the sampling distribution is approximately normal.
- (d) $\mu_{\bar{x}} = 50, \sigma_{\bar{x}} = 2$, the sampling distribution is approximately normal.

Answer: D)

Problem 5

An auto insurance company has 32,000 clients, and 5% of their clients submitted a claim in the past year. We will take a sample of 3,200 clients and determine how many of them have submitted a claim in the past year. What is the sampling distribution of \hat{p} ?

- a) $\hat{p} \sim N(3200, 0.2)$
- b) $\hat{p} \sim N(160, 152)$
- c) $\hat{p} \sim N(0.05, 0.003852)$
- d) Can not be determined

Answer: C) $\sigma_{\hat{p}} = \sqrt{\frac{0.05(1-0.05)}{3200}} = 0.00385$

Problem 6

A survey indicates that 60% of adults in a certain city support a new public transportation project. If a random sample of 200 adults is taken, what is the probability that the sample proportion supporting the project is greater than 65%?

- a) Approximately 0.0745
- b) Approximately 0.1151
- c) Approximately 0.2611
- d) Approximately 0.3821

Answer: A) $\sigma_{\hat{p}} = \sqrt{\frac{0.6(1-0.6)}{200}} = 0.0346$, by CLT $\hat{p} \sim N(0.6, 0.0346)$. Next, we find z-score of 0.65: $z_0 = (0.65-0.6)/0.0346 = 1.445$. $P(Z > 1.445) = 0.0742$

The primary interest of applying the CLT to sample proportion is to find the probability of an event defined by the sampling distribution of sample proportions.

1. Which Probability to Find?


- $P[p_0 < \hat{p} < p_1] = ?$
- $P[\hat{p} > p_0] = ?$
- $P[\hat{p} < p_0] = ?$

Given Value: p_0

2. Input Information

Population Proportion: p


Sample Size: n




Report bugs to C. Peng

Sampling Distribution of Sample Proportions

Given Information
 $p_0 = 0.65$



Standard Normal Distribution N(0,1)



$$Z = \frac{\hat{p} - p}{\sqrt{\frac{p(1-p)}{n}}}$$

Question: $P(\hat{p} > 0.65) = ?$

Solution: The answer is given in the following steps.

Step 1. Recall the Z-score Transformation

$$Z = \frac{\hat{p} - 0.6}{\sqrt{\frac{0.6 \times (1-0.6)}{200}}}$$

Step 2. Z-score for $p_0 = 0.65$ is given by

$$Z_0 = \frac{0.65 - 0.6}{\sqrt{\frac{0.6 \times (1-0.6)}{200}}} = 1.4434.$$

Step 3. Finding the left-tail probabilities

$$P(Z < 1.4434) = 0.9255.$$

Step 4. Note that

$$P(\hat{p} > 0.65) = P(Z > Z_1) = 1 - P(Z < 1.4434) = 1 - 0.9255 = 0.0745.$$

Step 5. That is, $P(\hat{p} > 0.65) = 0.0745$.

Problem 7

In a large population, 40% of individuals have a certain genetic trait. A random sample of 400 individuals is selected. Which of the following statements about the sampling distribution of the sample proportion is **incorrect**?

- a) The mean of the sampling distribution is 0.40.
- b) The standard deviation of the sampling distribution is approximately 0.0245.
- c) The shape of the sampling distribution is approximately normal.
- d) The standard deviation of the sampling distribution is larger than the standard deviation of the population.

Answer: D) Population standard deviation $\sqrt{0.4(1 - 0.4)} = \sqrt{0.24}$, while the standard deviation of the sampling distribution is $\sqrt{0.4(1 - 0.4)/400} = \sqrt{0.24}/20$. Therefore, the standard deviation of the sampling distribution is **smaller** than the standard deviation of the population.

Problem 8

A population has a mean of 50 and a standard deviation of 8. A random sample of size 64 is taken. Which of the following statements about the sampling distribution of the sample mean is **incorrect**?

- a) The mean of the sampling distribution is 50.
- b) The standard deviation of the sampling distribution is 1.

- c) The sampling distribution is approximately normal.
 d) The standard deviation of the sampling distribution is 8.

Answer: D) The standard deviation of the sampling distribution is $8/\sqrt{50} \approx 1.131$

Problem 9

A sample of size 49 is taken from a population with a mean of 200 and a standard deviation of 14. What is the probability that the sample mean will be less than 196?

- a) 0.0013
 b) 0.0228
 c) 0.0475
 d) 0.1587

Answer: B) Using the CLT, $\bar{X} \sim N\left(200, \frac{14}{\sqrt{49}}\right) = N(200, 2)$. $Z_0 = (196-200)/2 = -2 \rightarrow P(Z < -2) = 0.0228$

1. What to Find?

- Probability (P_0)
- Percentile (X_0)

2. Which Probability?

- $P[V_0 < \bar{X} < V_1] = ?$
- $P[\bar{X} > V_0] = ?$
- $P[\bar{X} < V_0] = ?$


Given Value: V_0

3. Input Information

Population Mean: μ

Population Standard Deviation: σ

Sample Size: n




Report bugs to C. Peng


Sampling Distribution of Sample Means

Given Information

$V_0 = 196$



Standard Normal Distribution N(0,1)



$$Z = \frac{X - \mu}{\sigma/\sqrt{n}}$$

Question: $P(\bar{X} < 196) = ?$

Solution: The answer is given in the following steps.

Step 1. Recall the Z-score transformation

$$Z = \frac{\bar{X} - 200}{14/\sqrt{49}}$$

Step 2. Z-scores for $V_0 = 196$ is given by

$$Z_0 = \frac{196 - 200}{14/\sqrt{49}} = -2.$$

Step 3. Note that

$$P(\bar{X} < 196) = P(Z < -2) = 0.0228.$$

Step 4. Therefore,

$$P(\bar{X} < 196) = 0.0228.$$

Problem 10

The lengths of pregnancies are normally distributed with a mean of 271 days and a standard deviation of 20 days. If 64 women are randomly selected, find the probability that they have a mean pregnancy between 271 days and 273 days.

- A). 0.2119
- B). 0.5517
- C). 0.2881
- D). 0.7881

Answer: C). First $\bar{X} \sim N\left(271, \frac{20}{\sqrt{64}}\right) = N(271, 2.5)$, Two corresponding z-scores are: $z_1 = (271-271)/2.5 = 0$, $z_2 = (273-271)/2.5 = 0.8$. $\rightarrow P(Z < 0.8) - P(Z < 0) = 0.7881 - 0.5 = 0.2881$

1. What to Find?

- Probability (P_0)
- Percentile (X_0)

2. Which Probability?

- $P[V_0 < \bar{X} < V_1] = ?$
- $P[\bar{X} > V_0] = ?$
- $P[\bar{X} < V_0] = ?$

Given Value #1: V_0


Given Value #2: V_1

3. Input Information

Population Mean: μ

Population Standard Deviation: σ


Sample Size: n




Report bugs to C. Peng

Sampling Distribution of Sample Means

Given Information
 $V_0 = 271$
 $V_1 = 273$



Standard Normal Distribution N(0,1)



$$Z = \frac{X - \mu}{\sigma/\sqrt{n}}$$

Question: $P(271 < \bar{X} < 273) = ?$

Solution: The answer is given in the following steps.

Step 1. Z-score Transformation

$$Z = \frac{\bar{X} - 271}{20/\sqrt{64}}$$

Step 2. Z-scores for $V_0=271$ and $V_1=273$ are given by

$$Z_0 = \frac{271 - 271}{20/\sqrt{64}} = 0,$$

$$Z_1 = \frac{273 - 271}{20/\sqrt{64}} = 0.8.$$

Step 3. Note that

$$P(271 < \bar{X} < 273) = P(0 < Z < 0.8)$$

$$= P(Z < 0.8) - P(Z < 0)$$

$$= 0.7881 - 0.5 = 0.2881.$$

Step 4. That is,

$$P(271 < \bar{X} < 273) = 0.2881.$$

Problem 11

Assume that the heights of women are normally distributed with a mean of 63.6 inches and a standard deviation of 2.5 inches. If 100 women are randomly selected, find the probability that they have a mean height greater than 63.0 inches.

- A). 0.2881
- B). 0.0082
- C). 0.9918
- D). 0.8989

Answer: C). First $\bar{X} \sim N\left(63.6, \frac{2.5}{\sqrt{100}}\right) = N(63.6, 0.25)$, the z-scores is: $z_1 = (63.0 - 63.6)/0.25 = -2.4$, $\rightarrow P(Z < -2.4) = 0.0082 \rightarrow P(\bar{X} > 63.0) = 1 - 0.0082 = 0.9918$.

1. What to Find?

- Probability (P_0)
- Percentile (X_0)

2. Which Probability?

- $P[V_0 < \bar{X} < V_1] = ?$
- $P[\bar{X} > V_0] = ?$
- $P[\bar{X} < V_0] = ?$


Given Value: V_0

3. Input Information

Population Mean: μ

Population Standard Deviation: σ

Sample Size: n




Report bugs to C. Peng


Sampling Distribution of Sample Means

Given Information

$V_0 = 63$



Standard Normal Distribution N(0,1)


$$Z = \frac{X - \mu}{\sigma/\sqrt{n}}$$

Question: $P(\bar{X} > 63) = ?$

Solution: The answer is given in the following steps.

Step 1. Z-score Transformation

$$Z = \frac{\bar{X} - 63.6}{2.5/\sqrt{100}}$$

Step 2. The Z-score for $V_0 = 63$ is

$$Z_0 = \frac{63 - 63.6}{2.5/\sqrt{100}} = -2.4$$

Step 3. Note that

$$P(Z < -2.4) = 0.0082$$

Step 4. Therefore,

$$\begin{aligned} P(\bar{X} > 63) &= P(Z > -2.4) \\ &= 1 - P(Z < -2.4) \\ &= 1 - 0.0082 = 0.9918 \end{aligned}$$

Problem 12

The body temperatures of adults are normally distributed with a mean of 98.6° F and a standard deviation of 0.60° F. If 36 adults are randomly selected, find the probability that their mean body temperature is greater than 98.4° F.

- A). 0.9360
- B). 0.0228
- C). 0.8188
- D). 0.9772

Answer D). First $\bar{X} \sim N\left(98.6, \frac{0.6}{\sqrt{36}}\right) = N(98.6, 0.1)$, the z-scores is: $z_1 = (98.4 - 98.6)/0.1 = -2$, $\rightarrow P(Z > -2) = 1 - P(Z < -2) = 1 - 0.02275 = 0.9772$.

1. What to Find?

- Probability (P_0)
- Percentile (X_0)

2. Which Probability?

- $P[V_0 < \bar{X} < V_1] = ?$
- $P[\bar{X} > V_0] = ?$
- $P[\bar{X} < V_0] = ?$


Given Value: V_0

3. Input Information

Population Mean: μ

Population Standard Deviation: σ

Sample Size: n




Report bugs to C. Peng

Sampling Distribution of Sample Means


Given Information

$V_0 = 98.4$



V_0

Standard Normal Distribution N(0,1)



z_0

$$Z = \frac{X - \mu}{\sigma/\sqrt{n}}$$

Question: $P(\bar{X} > 98.4) = ?$

Solution: The answer is given in the following steps.

Step 1. Z-score Transformation

$$Z = \frac{\bar{X} - 98.6}{0.6/\sqrt{36}}$$

Step 2. The Z-score for $V_0 = 98.4$ is

$$Z_0 = \frac{98.4 - 98.6}{0.6/\sqrt{36}} = -2$$

Step 3. Note that

$$P(Z < -2) = 0.0228$$

Step 4. Therefore,

$$P(\bar{X} > 98.4) = P(Z > -2)$$

$$= 1 - P(Z < -2)$$

$$= 1 - 0.0228 = 0.9772$$

Problem 13

A random sample of $n = 300$ measurements is drawn from a binomial population with probability of success .43. Give the mean and the standard deviation of the sampling distribution of the sample proportion, \hat{p} .

- A). $N(.57; .029)$
- B). $N(.43; .014)$
- C). $N(.57; .014)$
- D). $N(.43; .029)$

Answer D). $\sigma_{\hat{p}} = \sqrt{0.43(1 - 0.43)/300} \approx 0.0285$.

Problem 14.

The heights of adult women in a city are normally distributed with a mean of 165 cm and a standard deviation of 5 cm. What is the height that 95% of women are shorter than?

- a) 173.25 cm
- b) 171.45 cm
- c) 175.00 cm
- d) 174.60 cm

Answer: A.

1. What to Find?

- Probability (P_0)
- Percentile (X_0)

2. X_0 in Which Probability?


- $P[X_0 < X < V] = P_0$
- $P[V < X < X_0] = P_0$
- $P[X > X_0] = P_0$
- $P[X < X_0] = P_0$

Given Probability: P_0

3. Input Information

Population Mean: μ


Population Standard Deviation: σ



Report bugs to C. Peng

General Normal Distribution $N(\mu, \sigma)$


Given Information
 $P_0 = 0.95$



X_0

$$Z = \frac{X - \mu}{\sigma}$$

Standard Normal Distribution



Z_0

Question: Given $P(X < X_0) = 0.95$, what is X_0 ?

Solution: The answer is given in the following steps.

Step 1. Recall Z-score Transformations

$$Z = \frac{X - 165}{5} \text{ and } Z = \frac{X_0 - 165}{5}.$$

Step 2. The given condition $P(X < X_0) = 0.95$ is equivalent to

$$P(Z < Z_0) = 0.95$$

which gives, $Z_0 = 1.64$.

Step 3. Note that,

$$\frac{X_0 - 165}{5} = Z_0 = 1.64.$$

Step 4. Solve for X_0 from the above equation, we have,

$$X_0 = 165 + (1.64) \times 5 = 173.2.$$

Problem 15.

The test scores in a large class are normally distributed with a mean of 75 and a standard deviation of 10. What is the probability that a randomly selected student scored between 70 and 85?

- a) 0.5328
- b) 0.6826
- c) 0.7745
- d) 0.3413

Answer: A

1. What to Find?

- Probability (P_0)
- Percentile (X_0)

2. Which Probability?

- $P[V_0 < X < V_1] = ?$
- $P[X > V_0] = ?$
- $P[X < V_0] = ?$

Given Value #1: V_0

Given Value #2: V_1

3. Input Information

Population Mean: μ

Population Standard Deviation: σ

General Normal Distribution $N(\mu, \sigma)$

Given Information
 $V_0 = 70$
 $V_1 = 85$

Standard Normal Distribution

$$Z = \frac{X - \mu}{\sigma}$$

Question: $P(70 < X < 85) = ?$

Solution: The answer is given in the following steps.

Step 1. Recall Z-score Transformation

$$Z = \frac{X - 75}{10}$$

Step 2. Z-scores for $V_0 = 70$ and $V_1 = 85$ are given respectively by

$$Z_0 = \frac{70 - 75}{10} = -0.5 \text{ and } Z_1 = \frac{85 - 75}{10} = 1.$$

Step 3. The two left-tail Probabilities are

$$P(Z < 1) = 0.8413 \text{ and } P(Z < -0.5) = 0.3085.$$

Step 4. Note that


$$\begin{aligned} P(Z_0 < Z < Z_1) &= P(Z < Z_1) - P(Z < Z_0) \\ &= P(Z < 1) - P(Z < -0.5) \\ &= 0.8413 - 0.3085 = 0.5328. \end{aligned}$$

Summary of Weekly Assignment #5

The class boundary is: 30,40,50,60,70,80,90,100

cut.data.freq	Freq	midpts	rel.freq	cum.freq	rel.cum.freq
[3e+01,4e+01]	1	35.00	0.02	1	0.02
(4e+01,5e+01]	0	45.00	0.00	1	0.02
(5e+01,6e+01]	0	55.00	0.00	1	0.02
(6e+01,7e+01]	2	65.00	0.05	3	0.07
(7e+01,8e+01]	6	75.00	0.15	9	0.22
(8e+01,9e+01]	17	85.00	0.41	26	0.63
(9e+01,1e+02]	15	95.00	0.37	41	1.00

Probability Distribution Histogram



1. Five Number Summary :

The five-number summary is used to describe the shape of the distribution of a given numerical data. It consists of five numbers: minimum data value, first quartile, median, the third quartile, and the maximum data value.

The five-number summary of this given data set is:

stats	value
Min.	30.00
1st Qu.	85.00
Median	90.00
3rd Qu.	95.00
Max.	95.00

2. Boxplot :

The boxplot is a geometric representation of the five-number summary. The boxplot of the given data set is given below.

